

The oldest record of Alvarezsauridae (Dinosauria: Theropoda) in the Northern Hemisphere

Averianov A., Sues H.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2017, Public Library of Science. All rights reserved. This is an open access article, free of all copyright, and may be freely reproduced, distributed, transmitted, modified, built upon, or otherwise used by anyone for any lawful purpose. The work is made available under the Creative Commons CC0 public domain dedication. Procoelous caudal vertebrae, a carpometacarpus with a hypertrophied metacarpal II, and robust proximal and ungual phalanges of manual digit II of a small theropod dinosaur from the Upper Cretaceous (Turonian) Bissekty Formation at Dzharakuduk, Uzbekistan, show unequivocal synapomorphies of the clade Alvarezsauridae and thus are referred to it. The caudal vertebrae have a unique longitudinal canal within the neural arch. The carpometacarpus, with metacarpal III occupying about one third of the width of the carpometacarpus, shows the most plesiomorphic stage of the evolution of the forelimb among known alvarezsaurids. The proximal phalanx of manual digit II differs from the corresponding bone in Parvicursorinae in having a less asymmetrical proximal articular surface without a dorsal process and short ventral ridges. The ungual phalanx of manual digit II has laterally open ventral foramina. The Bissekty alvarezsaurid possibly represents a basal parvicursorine and is the stratigraphically oldest known alvarezsaurid in Asia known to date.

<http://dx.doi.org/10.1371/journal.pone.0186254>

References

- [1] Chiappe LM, Norell MA, Clark JM. The Cretaceous, short-armed Alvarezsauridae: *Mononykus* and its kin. In: Chiappe LM, Witmer LM, editors. *Mesozoic Birds: Above the Heads of Dinosaurs*. Berkeley: University of California Press; 2002. p. 87–120.
- [2] Novas FE. Alvarezsauridae, Cretaceous basal birds from Patagonia and Mongolia. *Memoirs of the Queensland Museum*. 1996; 39(3):675–702.
- [3] Xu X, Sullivan C, Pittman M, Choiniere JN, Hone D, Upchurch P, et al. A monodactyl nonavian dinosaur and the complex evolution of the alvarezsauroid hand. *Proceedings of the National Academy of Sciences USA*. 2011; 108(6):2338–42. <https://doi.org/10.1073/pnas.1011052108> PMID: 21262806
- [4] Longrich NR, Currie PJ. *Albertonykus borealis*, a new alvarezsaur (Dinosauria: Theropoda) from the early Maastrichtian of Alberta, Canada: implications for the systematics and ecology of the Alvarezsauridae. *Cretaceous Research*. 2009; 30(1):239–52.
- [5] Naish D, Dyke GJ. *Heptasteornis* was no ornithomimid, troodontid, dromaeosaurid or owl: the first alvarezsaurid (Dinosauria: Theropoda) from Europe. *Neues Jahrbuch für Geologie und Paläontologie Monatshefte*. 2004; 7:385–401.
- [6] Choiniere JN, Xu X, Clark JM, Forster CA, Guo Y, Han F. A basal alvarezsauroid theropod from the early Late Jurassic of Xinjiang, China. *Science*. 2010; 327(5965):571–4. <https://doi.org/10.1126/science.1182143> PMID: 20110503

- [7] Choiniere JN, Clark JM, Norell M, Xu X. Cranial osteology of *Haplocheirus sollers* Choiniere et al., 2010 (Theropoda, Alvarezsauridae). *American Museum Novitates*. 2014; 3816:1–44.
- [8] Novas FE. Anatomy of *Patagonykus puertai* (Theropoda, Avialae, Alvarezsauridae), from the Late Cretaceous of Patagonia. *Journal of Vertebrate Paleontology*. 1997; 17(1):137–66.
- [9] Averianov AO, Sues H-D. Correlation of Late Cretaceous continental vertebrate assemblages in Middle and Central Asia. *Journal of Stratigraphy*. 2012; 36(2):462–85.
- [10] Archibald JD, Sues H-D, Averianov AO, King C, Ward DJ, Tsaruk OI, et al. Précis of the Cretaceous paleontology, biostratigraphy and sedimentology at Dzharakuduk (Turonian?-Santonian), Kyzylkum Desert, Uzbekistan. *Bulletin of the New Mexico Museum of Natural History and Science*. 1998; 14:21–8.
- [11] Nessov LA, Archibald JD, Kielan-Jaworowska Z. Ungulate-like mammals from the Late Cretaceous of Uzbekistan and a phylogenetic analysis of Ungulatomorpha. *Bulletin of Carnegie Museum of Natural History*. 1998; 34:40–88.
- [12] Redman CM, Leighton LR. Multivariate faunal analysis of the Turonian Bissekty Formation: Variation in the degree of marine influence in temporally and spatially averaged fossil assemblages. *Palaios*. 2009; 24(1):18–26.
- [13] Gradstein FM, Ogg JG, Schmitz MD, Ogg GM. *A Geologic Time Scale 2012*. Amsterdam: Elsevier; 2012. xviii + 1144 p.
- [14] Xu X, Upchurch P, Ma Q, Pittman M, Choiniere JN, Sullivan C, et al. Osteology of the Late Cretaceous alvarezsaurid *Linhenykus monodactylus* from China and comments on alvarezsaurid biogeography. *Acta Palaeontologica Polonica*. 2013; 58(1):25–46.
- [15] Xu X, Clark JM, Mo J, Choiniere J, Forster CA, Erickson GM, et al. A Jurassic ceratosaur from China helps clarify avian digital homologies. *Nature*. 2009; 459:940–4. <https://doi.org/10.1038/nature08124> PMID: 19536256
- [16] Owen R. Report on British fossil reptiles. Part II. Report of the British Association for Advancement of Science. 1842:60–204.
- [17] Seeley HG. On the classification of the fossil animals commonly named Dinosauria. *Proceedings of the Royal Society of London*. 1887; 43(260):165–71.
- [18] Marsh OC. Classification of Dinosauria. *American Journal of Science, Series 3*. 1881; 23:81–6.
- [19] Gauthier JA. Saurischian monophyly and the origin of birds. *Memoirs of the California Academy of Sciences*. 1986; 8:1–55.
- [20] Bonaparte JF. Los vertebrados fósiles de la Formación Río Colorado, de la Ciudad de Neuquén y cercanías, Cretácico Superior, Argentina. *Revista del Museo Argentino de Ciencias Naturales, Paleontología*. 1991; 4(3):17–123.
- [21] Perle A, Norell MA, Chiappe LM, Clark JM. Flightless bird from the Cretaceous of Mongolia. *Nature*. 1993; 362:623–626.
- [22] Suzuki S, Chiappe LM, Dyke GJ, Watabe M, Barsbold R, Tsogtbaatar K. A new specimen of *Shuvuuia deserti* Chiappe et al., 1998 from the Mongolian Late Cretaceous with a discussion of the relationships of alvarezsaurids to other theropod dinosaurs. *Natural History Museum of Los Angeles County Contributions in Science*. 2002; 494:1–18.
- [23] Martinelli AG, Vera EI. *Achillesaurus manazzonei*, a new alvarezsaurid theropod (Dinosauria) from the Late Cretaceous Bajo de la Carpa Formation, Río Negro Province, Argentina. *Zootaxa*. 2007; 1582:1–17.
- [24] Chiappe LM, Coria RA. A new specimen of *Patagonykus puertai* (Theropoda: Alvarezsauridae) from the Late Cretaceous of Patagonia. *Ameghiniana*. 2003; 40(1):119–22.
- [25] Karhu AA, Rautian AS. A new family of Maniraptora (Dinosauria: Saurischia) from the Late Cretaceous of Mongolia. *Paleontological Journal*. 1996; 30(5):583–92.
- [26] Agnolin FL, Powell JE, Novas FE, Kundrát M. New alvarezsaurid (Dinosauria, Theropoda) from uppermost Cretaceous of north-western Patagonia with associated eggs. *Cretaceous Research*. 2012; 35:33–56.
- [27] Xu X, Wang D-Y, Sullivan C, Hone DWE, Han F, Yan R-H, et al. A basal parvicursorine (Theropoda: Alvarezsauridae) from the Upper Cretaceous of China. *Zootaxa*. 2010; 2413:1–19.
- [28] Hutchinson JR, Chiappe LM. The first known alvarezsaurid (Theropoda: Aves) from North America. *Journal of Vertebrate Paleontology*. 1998; 18(3):447–50.
- [29] Holtz TR, Jr. The arctometatarsalian pes, an unusual structure of the metatarsus of Cretaceous Theropoda (Dinosauria: Saurischia). *Journal of Vertebrate Paleontology*. 1994; 14(4):480–519.
- [30] Marsh OC. Notice of new reptiles from the Laramie Formation. *American Journal of Science, Series 3*. 1892; 43:449–53.
- [31] Salgado L, Coria RA, Arcucci AB, Chiappe LM. Restos de Alvarezsauridae (Theropoda, Coelurosauria) en la Formación Allen (Campaniano-Maastrichtiano), en Salitral Ojo de Agua, Provincia de Río Negro, Argentina. *Andean Geology*. 2009; 36(1):67–80.
- [32] Alifanov VR, Barsbold R. *Ceratonykus oculatus* gen. et sp. nov., a new dinosaur (?Theropoda, Alvarezsauria) from the Late Cretaceous of Mongolia. *Paleontological Journal*. 2009; 43(1):86–99.

- [33] Chiappe LM, Norell MA, Clark JM. The skull of a new relative of the stem-group bird *Mononykus*. *Nature*. 1998; 392:275-8.
- [34] Turner AH, Nesbitt SJ, Norell MA. A large alvarezsaurid from the Late Cretaceous of Mongolia. *American Museum Novitates*. 2009; 3648:1-14.
- [35] Hone DWE, Choiniere JN, Tan Q, Xu X. An articulated pes from a small parvicursorine alvarezsauroid dinosaur from Inner Mongolia, China. *Acta Palaeontologica Polonica*. 2013; 58(3):453-8.
- [36] Pittman M, Xu X, Stiegler JB. The taxonomy of a new parvicursorine alvarezsauroid specimen IVPP V20341 (Dinosauria: Theropoda) from the Upper Cretaceous Wulansuhai Formation of Bayan Mandahu, Inner Mongolia, China. *PeerJ*. 2015; 3:e986. <https://doi.org/10.7717/peerj.986> PMID: 26082871
- [37] Nesbitt SJ, Clarke JA, Turner AH, Norell MA. A small alvarezsaurid from the eastern Gobi Desert offers insight into evolutionary patterns in the Alvarezsauroidae. *Journal of Vertebrate Paleontology*. 2011; 31 (1):144-53.
- [38] Makovicky PJ, Apesteguía S, Gianechini FA. A new coelurosaurian theropod from the La Buitrera fossil locality of Río Negro, Argentina. *Fieldiana Life and Earth Sciences*. 2012; 5:90-8. <https://doi.org/10.3158/2158-552-5.1.90>